



ERECTILE DYSFUNCTION (IMPOTENCE)

**Information about your condition from
The British Association of Urological Surgeons (BAUS)**

You have been given this leaflet to provide you with detailed information about erectile dysfunction.

We have consulted specialist surgeons during its preparation, so it represents best practice in UK urology. You should use it in addition to any advice already given to you.

To view the online version of this leaflet, type the text below into your web browser:

[http://www.baus.org.uk/_userfiles/pages/files/Patients/Leaflets/Erectile dysfunction.pdf](http://www.baus.org.uk/_userfiles/pages/files/Patients/Leaflets/Erectile%20dysfunction.pdf)


Key Points

- Erectile dysfunction is a common problem in men over the age of 40
- Lifestyle modification such as increasing activity, stopping smoking, controlling high blood pressure and managing blood sugar levels are helpful in all patients
- Specific first-line treatment with tablets can be started by your general practitioner
- Some men are not eligible for free treatment of their erectile dysfunction under the NHS

What is the erectile dysfunction?

Erectile dysfunction (impotence) is the inability to get or keep an erection sufficient for sexual intercourse. One in ten men (10%) suffer from impotence and it is seen in almost one third (30%) of diabetic men.

The ability to get an erection is important to most men, old and young. However, many men suffer erectile dysfunction in silence, without seeking help or advice.



How do normal erections occur?

If he is to achieve an erection, a man needs:

- an adequate circulating level of hormones;
- an adequate blood flow to the penis;
- intact nerves supplying the penis; and
- an appropriate sexual desire.

If there is a problem with one or more of these mechanisms, erection may fail. During arousal, nerve impulses travel from the brain to the penis and trigger smooth muscle relaxation in the penis. This encourages blood to flow into the erectile tissues.

As the penis fills with blood, it enlarges and becomes erect. During enlargement, the veins in the penis become compressed, blocking the flow of blood out of the penis so that erection occurs. The penis remains erect until ejaculation is completed or sexual arousal ceases.

What physical causes are there for erectile dysfunction?

- **Hormone imbalance** – a deficiency of male hormones can reduce your desire or interest in sexual function
- **Nerve damage** - this can result in reduced sensitivity, or reduced signals to your penis to release the chemicals that cause
- **Disease of the blood vessels** - blood vessels often become narrowed and hardened with increasing age. This reduces blood supply to your penis. If the blood supply is poor, your penis may not fill with blood, the veins will not be compressed and what blood there is will leak out of the penis. Thus, your erection will not be maintained
- **Trauma** - such as injury to your spinal cord

- **Pelvic surgery** - some cancer operations on the prostate, bladder or bowel may result in nerve damage leading to erectile dysfunction
- **Drugs** - some drugs (especially those used to treat high blood pressure, depression and anxiety and recreational drugs) may cause erectile dysfunction
- **Smoking and alcohol** - those who smoke and drink are more likely to suffer from erectile dysfunction.

In men with diabetes, the commonest causes of erectile dysfunction are disease of the blood vessels and nerve damage (often in combination).

Can psychological problems cause erectile dysfunction?

Yes. It is very common to see a combination of psychological and physical causes, but pure psychological causes are seen in less than 1 in 10 (10%) of all affected patients. Erectile dysfunction can be caused by stress, depression, anxiety, relationship problems, embarrassment, guilt and other psychological issues.

When a man has difficulty getting an erection, whatever the cause, he will often experience pressure to perform. This can lead to a feeling of inadequacy and a sense of loss of manhood (called **performance anxiety**). These are all common emotions for men with erectile dysfunction.

What can I do about the problem?

You should talk about it with your partner, doctor or nurse. They will advise you about any tests which are needed.

There are some simple tests that your GP can do, such as blood tests to measure your lipids (cholesterol), blood sugar (diabetes check) and testosterone levels (hormone check), and measuring your blood pressure. Blood tests should be carried out first thing in the morning, preferably around 09.00 hr, and you should be starved from the night before.

If you can get any conditions (e.g. raised cholesterol, high blood pressure & diabetes) under control and get your weight to a healthy level, this is an effective starting point in treating your erectile dysfunction.

Once this assessment has been completed, other treatment options can be discussed with your GP.

What treatments are available?

It is up to you and your sexual partner to decide what treatment you choose. We will, of course, give you guidance on what we feel is most appropriate for you.

Specific treatment is only available on the NHS (Schedule 2) to patients who:

- have diabetes mellitus, multiple sclerosis, Parkinson's disease, poliomyelitis;
- have renal failure treated by dialysis or transplantation;
- have had radical pelvic surgery (e.g. radical prostatectomy) or have been treated for prostate cancer (using surgery and other treatments);
- have had severe pelvic injury, single-gene neurological disease, spinal cord injury or spina bifida; or
- are not included in the above categories but were receiving NHS treatment (such as Caverject™, Viagra™ or Viridal™) for their erectile dysfunction on or before 14 September 1998.

Originally, the Department of Health guidance stated that prescribing for those with erectile dysfunction causing “**severe distress**” should only be done by specialist services.

It is now possible for patients with severe distress due to erectile dysfunction to be prescribed generic sildenafil from their own GP, rather than having to attend a specialist service.


If you do not have any of the conditions listed above, you will need to get a private prescription from your GP. In summary, the basic treatment options are:

[Psychosexual counselling](#)

Some men need counselling and will be referred to a specialist in this area. Counselling can be part, or all, of the treatment required.

[Vacuum erection assistance devices](#)

Vacuum erection assistance devices are a non-invasive method of obtaining and sustaining an erection in men with impotence (erectile dysfunction).



To use the device, you insert your penis into a cylinder (above) with plenty of lubrication to provide a seal at its base. You get an erection by creating a negative pressure inside the cylinder, using a hand-operated or battery-powered pump.

Tablet treatment

First-line treatment for most patients is now tablet treatment using sildenafil (generic or Viagra™), tadalafil (Cialis™), vardenafil (Levitra™) or avanafil (Spedra™).

Your GP will prescribe the tablets for you in the first instance. An NHS prescription only allows you to be prescribed one tablet per week, but generic sildenafil is not subject to these restrictions.

Hormone treatment

This is only offered to patients who are deficient in male hormones. Medication can restore hormone imbalance and but does not always restore potency.

Hormone treatment will not improve erections in men who do not have hormone deficiency (and may even be harmful in this situation).

Self-injection therapy

This treatment involves injecting a drug into the side of your penis each time you want an erection.


The injection causes the muscles in your penis to relax which encourages blood to flow into your penis. If you choose this option, we will train you to inject yourself.

Injection therapy is very effective but some men find the idea difficult to accept. Injections can be used up to twice a week but you should never inject yourself more than once in any 24-hour period.

As with all drugs, there are some side-effects. For example, your erection may not go down and you then need to come to hospital to have it reversed. Fortunately, this only happens in 1 in 100 patients (1%).

Intra-urethral pellets or cream

This involves insertion of a pellet of prostaglandin into the urethra (waterpipe).



Only 35 to 40% of patients get good quality erections, and the pellet can cause pain or facial flushing.

[Penile implants](#)

This is an invasive surgical procedure which involves putting prostheses (implants) into your penis to allow you to achieve erections for sexual intercourse.

Penile implants are reserved for patients who have tried and failed other medical treatments such as tablets, injections, vacuum devices or pellets. It may also be used in patients with other conditions in which erections have been affected, such as following priapism (prolonged painful erections) or in men with Peyronie's disease.



The entire device is implanted into the body. It can be felt through the skin but is not visible on the outside.

What sources were used to prepare this leaflet?

This leaflet uses information from consensus panels and other evidence-based sources including:

- the [Department of Health \(England\)](#);
- the [Cochrane Collaboration](#); and
- the [National Institute for Health and Care Excellence \(NICE\)](#).

It also contains advice about the prescription of drugs. All NHS hospitals have local arrangements with their Clinical Commissioning Groups (CCGs) about which medicines can be used. Thus, some drugs mentioned cannot

be prescribed by local hospitals.

Your treatment will be planned with the doctors responsible for your care, considering not only which drugs are, or are not, available at your local hospital but also what is necessary to give you the best quality of care.

Healthcare professionals are advised to check prescribing arrangements with their local hospital or CCG.

This leaflet also follows style guidelines from:

- the [Royal National Institute for Blind People \(RNIB\)](#);
- the [Information Standard](#);
- the [Patient Information Forum](#); and
- the [Plain English Campaign](#).

Disclaimer

We have made every effort to give accurate information in this leaflet, but there may still be errors or omissions. BAUS cannot accept responsibility for any loss from action taken (or not taken) as a result of this information.

PLEASE NOTE

The staff at BAUS are not medically trained, and are unable to answer questions about the information provided in this leaflet. If you do have any questions, you should contact your urologist, specialist nurse or GP.